

08-OCT-2009  
2009-3372

GROUP	<b>1B</b>	INSECTICIDE
-------	-----------	-------------

**DIAZINON 5G  
INSECTICIDE**

**COMMERCIAL**

This product is not to be used in and around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.

**CAUTION                      POISON**

**READ THE LABEL BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN**

**REGISTRATION NO. 12538**

**PEST CONTROL PRODUCTS ACT**

**GUARANTEE: Diazinon 5%**

**Warning, contains the allergens soy and walnut shells.**

**UNITED AGRI PRODUCTS CANADA INC.**

**789 Donnybrook Drive**

**Dorchester, Ontario**

**N0L 1G5**

**1-800-265-4624**

**NET CONTENTS: 15 kg**

## **GENERAL PRECAUTIONS:**

Use only as recommended on the label. Do not reuse container. Apply by ground only. DO NOT APPLY BY AIR.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: [www.croplife.ca](http://www.croplife.ca)

## **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, please note that DIAZINON 5G Insecticide contains a Group 1B insecticide. Any insect population may contain individuals naturally resistant to DIAZINON 5G Insecticide and other Group 1B insecticide. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of DIAZINON 5G Insecticide or other Group 1B insecticides with different groups that control the same pathogens.
- Avoid application of more than the indicated number of applications of DIAZINON 5G Insecticide or other insecticides in the same group in a season.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting, record keeping and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information and to report suspected resistance, contact the Technical Service, United Agri Products, 1-800-265-4624 or at [www.uap.ca](http://www.uap.ca).

## **DIRECTIONS FOR USE:**

VEGETABLES: ROOT MAGGOT CONTROL on Broccoli & Brussels sprouts, Cabbage, and Cauliflower.

SEED BEDS: Apply 150 grams per 100 lineal meters of furrow.

TRANSPLANTING: Apply to soil within 10 cm of the plant at the rate of 80 gm per 10 lineal meters of row.

On RADISH: Apply 175 gm per 100 lineal meters in furrow at seeding time (45 kg per ha at 38 cm row spacing).

To control CARROT RUST FLY (maggot) on Carrots and Parsnips, apply 22 kg per ha in furrow (38 cm apart) at seeding time, (equivalent to 85 gm per 100 lineal meters).

To control CARROT RUST FLY (adult), use Diazinon 500 E according to the label. On sandy soil, maintain 0.6-1.2 cm of soil between the insecticide and seed.

To control CORN ROOTWORM LARVAE on Corn, apply 22 kg per ha in a 7.5-15 cm band over the row 1-2.5 cm above the seed and in front of the press wheel, but not in direct contact with the seed.

To control ONION MAGGOT LARVAE on Onions and Green onions, apply 22 kg per ha (up to 45 kg in organic soil) to furrows at planting time or seeding time (equivalent to 85 gm/180 gm per 100 lineal meters).

To control VINEGAR FLIES on Tomatoes, apply 11-22 kg per ha. Broadcast uniformly over the soil surface when flies first appear. Repeat at 7-10 day intervals or as required. May be applied up to the day of harvest.

To control ROOT MAGGOTS on Rutabagas (turnips), apply 55 kg per ha (or 40 gm per 10 lineal meters of row 70 cm apart) in furrow at seeding time. Make second application two weeks after thinning. Use with caution on sandy soils as injury and reduced stand may result. To control adult flies, use Diazinon 500 E.C. according to the label.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. Do not use on humans, pets, livestock. May be fatal if swallowed. Do not get into eyes. Avoid contact with skin and inhalation of dust. Wear NIOSH approved respirator. Wash thoroughly after handling and before eating, drinking and smoking. Avoid contamination of food, feed or food utensils. When applying wear coveralls over work clothes and chemical resistant gloves during loading, application, irrigation and clean-up activities.

SYMPTOMS OF POISONING: Headaches, giddiness, blurred vision, nervousness, weakness, nausea, cramps, diarrhea, discomfort in the chest, sweating, miosis, tearing, salivation and other excessive respiratory tract secretion, vomiting, cyanosis, papilledema, uncontrollable muscle twitches, convulsions, coma, loss of sphincter control and loss of reflexes.

FISH AND WILDLIFE CAUTION: This product is toxic to birds, fish and wildlife, including waterfowl. Do not contaminate domestic or irrigation water supplies, lakes, streams and ponds. Birds and waterfowl feeding or drinking on treated area may

be killed. Do not apply where there are resident populations of waterfowl or where waterfowl may be expected to inhabit the area. Do not exceed label rates because rates above those recommended significantly increase hazards to birds and waterfowl. Avoid overlapping. Irrigation (watering) is recommended, applying a minimum of 0.7 cm (0.25 inches) of water immediately after application to reduce the hazards to birds. Stop irrigation as soon as puddling occurs.

TOXIC TO BEES.

FIRST AID: Take container, label or product name and Pest Control Product Registration number with you when seeking medical attention.

IF SWALLOWED, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION: Diazinon is an organophosphate insecticide and, as such, a cholinesterase inhibitor. If cholinergic systems occur, give atropine as an antidote followed by Pralidoxime (2-PAM) and supporting therapy. Use 1 to 2 mg of atropine every 15 to 30 minutes if necessary.

DISPOSAL:

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
  2. Make the empty container unsuitable for further use.
  3. Dispose of the container in accordance with provincial requirements.
  4. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency.
- Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.