

2007-8095
22-Jan-2010
[Carton Label]



GROUP	2	HERBICIDE
-------	---	-----------

NIMBLE™
herbicide

A Dry Flowable 75% Herbicide

FOR SALE FOR USE ON WHEAT (SPRING, WINTER OR DURUM), SPRING BARLEY AND OATS, AND SEEDLING AND ESTABLISHED GRASS SPECIES FOR FORAGE AND SEED PRODUCTION ONLY

WETTABLE GRANULES

AGRICULTURAL

GUARANTEE: Thifensulfuron-methyl 50%
Tribenuron-methyl 25%

REGISTRATION NO.: **29467**
PEST CONTROL PRODUCTS ACT

WARNING: CONTAINS THE ALLERGENS MILK AND SULFITES

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

<p>IN CASE OF A MEDICAL EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-866-303-6950</p>
--

Cheminova Canada Inc.
22499 Jefferies Road, Unit C2,
Kilworth, Ontario N0L 1R0
Canada
1-888-316-6260

NET CONTENTS:

NIMBLE™ is a trademark of Cheminova

PRECAUTIONS:**• KEEP OUT OF REACH OF CHILDREN.**

- Avoid contact with skin, eyes and clothing. Wear goggles or face shield, coveralls, and suitable chemical resistant gloves during mixing and loading.
- Do not contaminate any body of water.
- May be harmful if swallowed, inhaled or absorbed through skin.
- Causes eye and skin irritation.
- When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.
- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before re-use.
- Do not enter or allow worker re-entry treated fields for 12 hours after treatment.
- DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.
- Do not apply by air.

NOTE: If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. For medical emergencies call 1-866-303-6950.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

[Booklet]



GROUP

2

HERBICIDE

NIMBLE™

herbicide

A Dry Flowable 75% Herbicide

FOR SALE FOR USE ON WHEAT (SPRING, WINTER OR DURUM), SPRING BARLEY AND OATS, AND SEEDLING AND ESTABLISHED GRASS SPECIES FOR FORAGE AND SEED PRODUCTION ONLY

WETTABLE GRANULES

AGRICULTURAL

GUARANTEE: Thifensulfuron-methyl 50%
Tribenuron-methyl 25%

REGISTRATION NO.: **29467**
PEST CONTROL PRODUCTS ACT

WARNING: CONTAINS THE ALLERGENS MILK AND SULFITES

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.

**IN CASE OF A MEDICAL EMERGENCY, CALL TOLL FREE,
DAY OR NIGHT: 1-866-303-6950**

Cheminova Canada Inc.
22499 Jefferies Road, Unit C2,
Kilworth, Ontario N0L 1R0
Canada
1-888-316-6260

NET CONTENTS:

NIMBLE™ is a trademark of Cheminova

PRECAUTIONS:**• KEEP OUT OF REACH OF CHILDREN.**

- Avoid contact with skin, eyes and clothing. Wear goggles or face shield, coveralls, and suitable chemical resistant gloves during mixing and loading.
- Do not contaminate any body of water.
- May be harmful if swallowed, inhaled or absorbed through skin.
- Causes eye and skin irritation.
- When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.
- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before re-use.
- Do not enter or allow worker re-entry treated fields for 12 hours after treatment.
- DO NOT SELL WATER SOLUBLE BAGS SEPARATELY.
- Do not apply by air.

NOTE: If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. For medical emergencies call 1-866-303-6950.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS:

- Overspray or drift to important wildlife habits such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided. Leave a 15-meter buffer zone between the last spray swath and the edge of any of these habitats.
- Do not use on highly variable soils that have large gravelly or sand areas, eroded knolls, or calcium deposits.
- Do not remove soil from treated fields for use on lawns, in gardens or for back fill.
- Do not contaminate irrigation water. Do not apply to land where tail water will be used to irrigate cropland.
- Do not apply more than 20 g per hectare of **NIMBLE herbicide** per crop year.

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following: Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, recreational grounds or similar

areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops.

DO NOT APPLY BY AIR.

Carefully observe sprayer clean-up instructions, as spray tank residues may damage crops other than wheat, barley or oats.

APPLY WITH A SURFACTANT REGISTERED FOR USE WITH **NIMBLE herbicide**.

GENERAL INFORMATION

NIMBLE herbicide is recommended for selective postemergence control or suppression of broadleaf weeds in wheat (spring, winter or durum), spring barley and oats not underseeded to legumes or grasses. **NIMBLE herbicide** is a 75% active ingredient herbicide formulated as a dry flowable dispersible granule. **NIMBLE herbicide** is to be mixed with water and a recommended surfactant and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile and does not freeze.

NIMBLE herbicide MUST be applied with a recommended surfactant, early postemergence, to the main flush of actively growing broadleaf weeds. Recommended surfactants are AGRAL* 90, AG-SURF*, and CITOWETT* PLUS.

Warm, moist growing conditions promote active weed growth and enhance the activity of **NIMBLE herbicide** by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details. **NIMBLE herbicide** stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

Canola, flax, lentils and alfalfa may be planted two months after an application of **NIMBLE herbicide**. Any crop may be planted in the year following the use of **NIMBLE herbicide**. One 320 g container of **Nimble herbicide** treats 16 ha. Each 80-gram bag treats 4 ha.

IMPORTANT INFORMATION: READ BEFORE USING

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

- Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- Do not use on lawns, walks, driveways, recreational grounds or similar areas.
- Prevent drift of spray to desirable plants.
- Do not contaminate any body of water, including irrigation water that may be used on other crops.
- Carefully observe sprayer clean-up instructions, as spray tank residues may damage crops other than wheat, spring barley or oats.

REPLANTING TO OTHER CROPS:

Minimum Crop Rotation Guidelines:

Minimum Interval	Crop
2 months	Canola, Flax, Lentils, Alfalfa
Next crop year	All crops

GRAZING/HARVEST RESTRICTIONS:

Wheat, spring barley or oats may not be grazed or fed to livestock within 7 days of application.

VARIETY RESTRICTIONS:

Limit use of **NIMBLE herbicide** to currently licensed varieties of wheat (spring, durum, winter), spring barley and oats. Consult Cheminova Canada, Inc., prior to use on unlicensed varieties for crop tolerance information. **DO NOT USE ON LEGER SPRING BARLEY OR BELVEDER WHEAT VARIETIES.**

APPLICATION INFORMATION:

APPLICATION PRECAUTIONS:

- DO NOT APPLY BY AIR.
- Do not apply to irrigated land where tail water will be used to irrigate cropland.
- Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls or calcium deposits.
- Overspray or drift to important wildlife habitats such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided. Leave a 15-meter buffer zone between the last spray swath and the edge of any of these habitats.
- Apply with an adjuvant registered for use with **NIMBLE Herbicide**.
- Use only water as a carrier. Other carriers may accelerate breakdown of **NIMBLE herbicide** and reduce its effectiveness.
- Do not apply to crops underseeded to legumes or grasses as injury to forage may result.
- Application to crops that are stressed by severe weather conditions, frost, low fertility, water logged soil (soil at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury.
- Under certain conditions, such as heavy rainfall, prolonged cool weather, frost or wide fluctuations in day/night temperatures, temporary lightening in crop colour, and occasionally, a slight reduction in crop height may occur.
- Do not apply more than 20 grams/hectare of **NIMBLE herbicide** per year.

APPLICATION INSTRUCTIONS:

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh or larger filter screens (metal or nylon). A spray pressure of 210-275 kPa is recommended.

Continuous agitation is required to keep **NIMBLE herbicide** in suspension. Apply uniformly. Avoid overlapping and shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

DIRECTIONS FOR USE

WHEAT (SPRING, DURUM, WINTER), SPRING BARLEY AND OATS

NIMBLE herbicide	
LOCATION: Not restricted	
CROP(S): wheat (spring, durum, winter), spring barley and oats only	
TIMING: Post-emergent application from 2-leaf to flag leaf (shot blade) growth stage of crop, prior to head emergence, but before the crop canopy will prevent thorough coverage of target weeds. For best results apply to young, actively growing weeds that are less than 10 cm in height or diameter. Thorough coverage of target weeds is essential.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare

Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS	2 litres/1000 litres of spray solution
MIXING: Mixing order is NIMBLE herbicide, then adjuvant. Refer to MIXING DIRECTIONS section for general instructions	
WEEDS CONTROLLED	
annual smartweed (green smartweed, lady's thumb)/ball mustard/chickweed ¹ /corn spurry/cow cockle/flixweed/groundsel (common)/hemp-nettle/kochia/lamb's quarters/narrow-leaved hawk's beard/redroot pigweed/Russian thistle/shepherd's purse/stinkweed/tartary buckwheat/volunteer rapeseed ² /volunteer sunflower/wild buckwheat ³ /wild mustard	
Notes	
¹ Apply when chickweed is small (1-6 leaf) and actively growing, but before crop canopy prevents thorough herbicide coverage of weeds. Chickweed emerging after application will not be controlled.	
² NIMBLE herbicide alone will not control imazamox and imazethapyr tolerant canola (e.g., Clearfield* canola).	
³ Apply to actively growing wild buckwheat in the cotyledon 1-3 leaf stage. Under environmental stress, such as dry weather, control may be reduced. Large plants may re-grow after treatment.	
WEEDS SUPPRESSED	
Canada thistle ⁴ /cleavers (1-3 whorls)/round leaved mallow ⁵ /scentless chamomile/Sow thistle ⁴ /stork's bill (2-6 leaf)/toadflax ⁶	
Notes	
⁴ Apply when the majority of thistles have emerged and are actively growing. For best top growth control, apply before bud stage when thistles are no larger than 15 cm in height and before crop canopy prevents thorough coverage of weeds. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be suppressed	
⁵ Apply to actively growing round leaved mallow from the 2-6 leaf stage (10-12 cm in height). NIMBLE herbicide will keep mallow stunted, but may not reduce overall populations.	
⁶ Apply to toadflax that is no larger than 15 cm in height. A control program for this weed includes both frequent tillage and chemical application.	
NOTE: Rainfall within 4 hours of application of NIMBLE herbicide will result in reduced weed control.	

NIMBLE herbicide + MCPA	
LOCATION: Not restricted	
CROP(S): wheat (spring, durum, winter), spring barley and oats only	
TIMING: Post-emergent application from full 3-leaf to just before flag leaf (shot blade) growth stage of crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
MCPA Amine or Ester 500	0.7-1.1 litres/hectare (350-560 grams acid equivalent/hectare)
OR	
MCPA Ester 500	0.56 litres/hectare (280 grams acid equivalent/hectare)
Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS	2 litres/1000 litres of spray solution
MIXING: Mixing order is NIMBLE herbicide , then MCPA to the tank while maintaining agitation, then adjuvant. Antifoaming agent may be required. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED BY NIMBLE herbicide+ MCPA AT 0.7-1.1 L/HA	

annual smartweed (green smartweed, lady's thumb)/ball mustard/burdocks (seedling)/chickweed¹/cocklebur²/corn spurry/cow cockle/false ragweed²/flixweed/giant ragweed²/hare's-ear mustard/hemp-nettle/Indian mustard/kochia²/lamb's quarters/plantain (common)/prickly lettuce/ragweed²/redroot pigweed/Russian pigweed²/Russian thistle/shepherd's purse/stinkweed/tartary buckwheat/tumble mustard/volunteer rapeseed/wild buckwheat³/wild mustard/wild radish/wormseed mustard

WEEDS CONTROLLED NIMBLE herbicide+ MCPA ESTER 500 AT 0.56 L/HA

Weeds controlled by **Nimble herbicide** as well as volunteer imazamox and imazethapyr tolerant canola(e.g., Clearfield* canola)(2-4 leaf stage)

Notes

¹ Apply when chickweed is small (1-6 leaf) and actively growing, but before crop canopy prevents thorough herbicide coverage of weeds. Chickweed emerging after application will not be controlled.

² Use minimum of 400 grams MCPA active ingredient per hectare.

³ Apply to actively growing wild buckwheat in the cotyledon 1-3 leaf stage. Under environmental stress, such as dry weather, control may be reduced. Large plants may re-grow after treatment.

WEEDS SUPPRESSED

Canada thistle⁴/round leaved mallow⁵/scentless chamomile/sow thistle⁴/stork's bill (2-6 leaf) (MCPA Ester only)

Notes

⁴ Apply when the majority of thistles have emerged and are actively growing. For best top growth control, apply before bud stage when thistles are no larger than 15 cm in height and before crop canopy prevents thorough coverage of weeds. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be suppressed.

⁵ Apply to actively growing round leaved mallow from the 2-6 leaf stage (10-12 cm in height). Nimble herbicide + MCPA will keep mallow stunted, but may not reduce overall populations.

NOTES: Ensure that NIMBLE herbicide is completely in suspension before adding MCPA. Consult the MCPA label for additional application instructions and use precautions.

WHEAT (SPRING, DURUM, WINTER) AND SPRING BARLEY

NIMBLE herbicide + 2,4-D AMINE or ESTER	
LOCATION: Not restricted	
CROP(S): wheat (spring, durum, winter) and spring barley only. DO NOT USE ON OATS	
TIMING: Post-emergent application from full 3-leaf to just before flag leaf (shot blade) growth stage of crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
Choose one of the following:	
1) 2,4-D Amine 500	0.84-1.1 litres/hectare
2) 2,4-D Ester LV700	0.6-0.8 litres/hectare
3) 2,4-D Ester LV600	

4) Acid equivalent. =420 – 550 gms active/ha.	0.7-0.9 litres/hectare
Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS	2 litres/1000 litres of spray solution
MIXING: Mixing order is NIMBLE herbicide , then 2,4-D to the tank while maintaining agitation, then adjuvant. Antifoaming agent may be required. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
annual smartweed (green smartweed, lady's thumb)/ball mustard/burdocks (seedling)/chickweed ¹ /cocklebur/common ragweed/corn spurry/cow cockle/false ragweed/flixweed/giant ragweed/hare's-ear mustard/hemp-nettle/Indian mustard/kochia/lamb's quarters/plantain (common)/prickly lettuce/redroot pigweed/Russian pigweed/Russian thistle/shepherd's purse/stinkweed/stork's bill (2-6 leaf) (2,4-D Ester only)/tartary buckwheat/tumble mustard/volunteer rapeseed ² /wild buckwheat ³ /wild mustard/wild radish/wormseed mustard	
Notes	
¹ Apply when chickweed is small (1-6 leaf) and actively growing, but before crop canopy prevents thorough herbicide coverage of weeds. Chickweed emerging after application will not be controlled.	
² For imazamox and imazethapyr tolerant canola (e.g., Clearfield* canola) consult the 2,4-D label for appropriate use rates.	
³ Apply to actively growing wild buckwheat in the cotyledon 1-3 leaf stage. Under environmental stress, such as dry weather, control may be reduced. Large plants may re-grow after treatment.	
WEEDS SUPPRESSED	
Canada thistle ⁴ /round leaved mallow ⁵ /scentless chamomile/sow thistle ⁴ /toadflax (2,4-D Ester only)	
Notes	
⁴ Apply when the majority of thistles have emerged and are actively growing. For best top growth control, apply before bud stage when thistles are no larger than 15 cm in height and before crop canopy prevents thorough coverage of weeds. NIMBLE herbicide+ 2,4-D will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be suppressed.	
⁵ Apply to actively growing round leaved mallow from the 2-6 leaf stage (10-12 cm in height). NIMBLE herbicide + 2,4-D will keep mallow stunted, but may not reduce overall populations. NOTES: Ensure that NIMBLE herbicide is completely in suspension before adding 2,4-D. Consult the 2,4-D label for additional application instructions and use precautions.	

SPRING WHEAT (INCLUDING DURUM) AND SPRING BARLEY

NIMBLE herbicide + ASSERT*	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley only	
TIMING: Post-emergent application from 2-leaf to 6-leaf growth stage of crop. Apply to wild oats from the 1-4 leaf stage.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare

ASSERT*	1.3 litres/hectare for 1-3 leaf stage of wild oats 1.6 litres/hectare for 4-leaf stage of wild oats
No adjuvant required.	
MIXING: Mixing order is NIMBLE herbicide then Spray Water pH Adjuster then Assert*.	
Ensure that NIMBLE herbicide is completely in suspension before adding pH Adjuster and that pH Adjuster is completely dissolved before adding Assert* while maintaining agitation. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled by NIMBLE herbicide alone, as well as wild oats.	
NOTES: Use spray preparation within 12 hours or product degradation may occur. Consult the Assert* label for additional application instructions, use precautions and recropping information.	

NIMBLE herbicide + ASSERT* + MCPA Ester 500	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley only	
TIMING: Post-emergent application from 4-leaf to 6-leaf growth stage of crop. Apply to wild oats from the 1-4 leaf stage.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
ASSERT*	BROWN AND DARK BROWN SOIL ZONES 1.3 litres/hectare for 1-3 leaf stage of wild oats 1.6 litres/hectare for 4-leaf stage of wild oats BLACK AND GREY WOODDED SOIL ZONES 1.6 litres/hectare
MCPA Ester 500	0.7-1.1 litres/hectare (350-560 grams acid equivalent/hectare)
No adjuvant required.	
MIXING: Mixing order is NIMBLE herbicide then Spray Water pH Adjuster then Assert* then MCPA Ester 500. Ensure that NIMBLE herbicide is completely in suspension before adding pH Adjuster and that pH Adjuster is completely dissolved before adding Assert*, while maintaining agitation. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled by NIMBLE herbicide alone, as well as wild oats.	
NOTES: Use spray preparation within 12 hours or product degradation may occur. Consult the Assert* and MCPA Ester 500 labels for additional application instructions, use precautions and recropping information.	

NIMBLE herbicide + AXIAL* 100EC Herbicide	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley only	
TIMING: Post-emergent application from 1 to flag leaf. For annual grass weeds, apply from 1 to 6-leaf stage, up to emergence of 4th tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
AXIAL* 100 EC Herbicide	400 mL/hectare for Persian dandelion only 600mL/hectare for wild oats, volunteer oats,

	green foxtail, yellow foxtail, volunteer canary seed, and proso millet.
ADIGOR* Adjuvant Or MERGE* Adjuvant	700 mL/hectare
MIXING: Mixing order is NIMBLE herbicide then Axial* 100 EC Herbicide, then Adigor* or Merge*, while maintaining agitation. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, plus wild oats, volunteer oats, green foxtail, yellow foxtail, Persian darnel, volunteer canary seed, and proso millet.	
NOTES: Do not harvest within 60 days for grain and straw and 30 days for hay. Do not graze treated crop for 7 days after application. Consult AXIAL* 100 EC Herbicide label for additional information on appropriate weed staging, application instructions, environmental and use precautions.	

NIMBLE herbicide + AXIAL* 100EC Herbicide + MCPA Ester 500	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley only	
TIMING: Post-emergent application from 1 to flag leaf. For annual grass weeds, apply from 1 to 6-leaf stage, up to emergence of 4 th tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
AXIAL* 100 EC Herbicide	400 mL/hectare for Persian darnel only 600mL/hectare for wild oats, volunteer oats, green foxtail, yellow foxtail, volunteer canary seed, and proso millet.
MCPA Ester 500	560 mL/hectare for volunteer imazamox and imazethapyr tolerant canola(e.g. Clearfield canola)(2 4 leaf stage). 700 mL/hectare for weeds controlled by MCPA Ester at this rate
ADIGOR* Adjuvant OR MERGE* Adjuvant	700 mL/hectare
MIXING: Mixing order is NIMBLE herbicide then Axial* 100 EC Herbicide, then MCPA Ester 500, and finally Adigor* or Merge*, while maintaining agitation. Refer to MIXING DIRECTIONS section for general instructions	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as weeds controlled by MCPA Ester 500 at 700 mL/hectare, or volunteer imazamox and imazethapyr tolerant canola (e.g. Clearfield* canola)(2-4 leaf stage) when applied at 560 mL/hectare, plus wild oats, volunteer oats, green foxtail, yellow foxtail, Persian darnel, volunteer canary seed, and proso millet.	
NOTES: Do not harvest within 60 days for grain and straw and 30 days for hay. Do not graze treated crop for 7 days after application. Consult AXIAL* 100 EC Herbicide and MCPA Ester labels for additional information on appropriate weed staging, application instructions, environmental and use precautions.	

NIMBLE herbicide + PUMA* SUPER	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley only	

TIMING: Post-emergent application from 1 to 6-leaf plus 3 tillers growth stage of spring and durum wheat; 1 to 5-leaf plus 2 tillers growth stage of spring barley. For annual grass weeds, apply from 1 to 6-leaf stage, up to emergence of third tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
PUMA* SUPER	0.5-1.0 litre/hectare for green foxtail 1.0 litre/hectare for wild oats and barnyard grass
No adjuvant required.	
MIXING: Mixing order is NIMBLE herbicide then Puma* Super, while maintaining agitation. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as green foxtail, barnyard grass and wild oats.	
NOTES: Consult the Puma* Super label for additional information on appropriate weed staging, application instructions, environmental and use precautions.	

NIMBLE herbicide + MCPA ESTER 500 + PUMA* SUPER	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley only	
TIMING: Post-emergent application from 1 to 6-leaf plus 3 tillers growth stage of spring and durum wheat; 1 to 5-leaf plus 2 tillers growth stage of spring barley. For annual grass weeds, apply from 1 to 6-leaf stage, up to emergence of third tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
MCPA ESTER 500	0.84 litres/hectare
PUMA* SUPER	0.5-1.0 litre/hectare for green foxtail 1.0 litre/hectare for wild oats and barnyard grass
No adjuvant required.	
MIXING: Mixing order is NIMBLE herbicide, then Puma* Super, while maintaining agitation, then MCPA Ester 500. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide + MCPA Ester 500 (0.84 litres/hectare) as well as green foxtail, barnyard grass and wild oats.	
NOTES: Consult the MCPA Ester 500 and Puma* Super labels for additional information on appropriate weed staging, application instructions, environmental and use precautions.	

NIMBLE herbicide + PUMA* 120 SUPER	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley only	
TIMING: Post-emergent application from 1 to 6-leaf plus 3 tillers growth stage of spring and durum wheat; 1 to 5-leaf plus 2 tillers growth stage of spring barley. For annual grass weeds, apply from 1 to 6-leaf stage, up to emergence of third tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
PUMA* 120 SUPER	385 mL/hectare for green foxtail only 770mL/hectare for wild oats, green foxtail, yellow foxtail and barnyard grass
No adjuvant required	
MIXING: Mixing order is NIMBLE herbicide then Puma* 120 Super, while maintaining agitation.	

Refer to MIXING DIRECTIONS section for general instructions.
WEEDS CONTROLLED
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as green foxtail, yellow foxtail, barnyard grass and wild oats.
NOTES: Consult the Puma* 120 Super label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

NIMBLE herbicide + PUMA* 120 SUPER + MCPA Ester 500	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley only	
TIMING: Post-emergent application from 1 to 6-leaf plus 3 tillers growth stage of spring and durum wheat; 1 to 5-leaf plus 2 tillers growth stage of spring barley. For annual grass weeds, apply from 1 to 6-leaf stage, up to emergence of third tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
PUMA* 120 SUPER	385 mL/hectare for green foxtail only 770mL/hectare for wild oats, green foxtail, yellow foxtail and barnyard grass
MCPA Ester 500	560 mL/hectare for volunteer imazamox and imazethapyr tolerant canola(e.g., Clearfield canola)(2-4 leaf stage). 840 mL/hectare for weeds controlled by MCPA Ester 500 at this rate.
No adjuvant required	
MIXING: Mixing order is NIMBLE herbicide then Puma* 120 Super, then MCPA Ester 500 while maintaining agitation. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone: green foxtail, yellow foxtail, barnyard grass and wild oats as well as weeds controlled by MCPA Ester 500 at 840 mL/hectare or volunteer imazamox and imazethapyr tolerant canola(e.g., Clearfield* canola)(2-4 leaf stage) when MCPA Ester 500 is applied at 560 mL/hectare.	
NOTES: Consult the MCPA Ester 500 and Puma* 120 Super labels for additional information on appropriate weed staging, application instructions, environmental and use precautions.	

NIMBLE herbicide + ACCORD*	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley(grown for feed purposes only) NOT UNDERSEEDED TO FORAGES	
TIMING: Post-emergent application from 2 to 5-leaf growth stage of the crop. See below for weed staging.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
ACCORD*	135 grams/hectare for most weeds listed below 165 grams/hectare for green foxtail 67 grams/hectare for cleavers and annual sowthistle
Adjuvant: MERGE* only	10 litres/1000 litres of spray solution (1.0% v/v)
MIXING: Mixing order is NIMBLE herbicide then Accord* then Merge* adjuvant. DO NOT USE ADDITIONAL NON-IONIC SURFACTANT. Refer to MIXING DIRECTIONS section for general instructions.	

WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as:	
green foxtail.....	1-5 leaf , 2 tillers
barnyard grass	1-5 leaf
volunteer flax	1-8 cm
cleavers	1-4 whorls
NOTES: Do not apply this tank mixture more often than once every 2 years. Do not graze the treated crops or cut for hay within 77 days of application. Consult the Accord* label for rotational cropping restrictions, application instructions, use precautions and other restrictions. Refer to the Accord* label for additional information on appropriate weed staging at application. Thorough coverage of foliage is important for consistent weed control.	

NIMBLE herbicide + ACCORD* + Puma* 120 Super	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) and spring barley(grown for feed purposes only) NOT UNDERSEEDED TO FORAGES	
TIMING: Post-emergent application from 2 to 5-leaf growth stage of spring and durum wheat or to barley at the 2 to 4-leaf stage. See below for weed staging.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
ACCORD*	67 grams/hectare for cleavers only
Puma* 120 Super	0.77 litres per hectare
MIXING: Mixing order is NIMBLE herbicide then Accord* then Puma* 120 Super. Merge adjuvant is NOT required in this tank mixture. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as: Wild oats, green foxtail, barnyard grass – 1 - 6-leaf stage up to emergence of third tiller Cleavers 1-4 whorls	
NOTES: Do not apply this tank mixture more often than once every 2 years. Do not graze the treated crops or cut for hay within 77 days (spring wheat, durum) or 80 days (spring barley) of application. Consult the Accord* label for rotational cropping restrictions, application instructions, use precautions and other restrictions. Refer to the Accord* and Puma* 120 Super labels for additional information on appropriate weed staging at application. Thorough coverage of foliage is important for consistent weed control	

SPRING WHEAT (EXCLUDING DURUM) AND SPRING BARLEY

NIMBLE herbicide + ATTAIN™ HERBICIDE TANK MIX		
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.		
CROP(S): spring wheat (excluding durum) and spring barley only, not underseeded to legumes		
TIMING: Post-emergent application from the 4-leaf to flag-leaf stage of spring wheat or spring barley and seedling stage to 4-leaf/whorls of weed.		
TANK MIX COMPONENTS AND APPLICATION RATES		
NIMBLE herbicide		20 grams/hectare
Attain* Herbicide	ATTAIN* A (fluroxypur)	Attain* A - 0.30 L/ha
Tank Mix	ATTAIN* B (2,4-D Ester)	Attain* B - 0.50 L/ha
Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS		2L/1000 litres of spray solution
MIXING: Mixing order is NIMBLE herbicide , then Attain* A, then Attain* B then surfactant. Refer to MIXING DIRECTIONS section for general instructions. Use a spray volume of 100 L/ha.		

WEEDS CONTROLLED
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as cleavers (excluding Group 2 resistant cleavers).
NOTES: Do not permit lactating dairy animals to graze fields within 7 days after application. Do not harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days prior to slaughter. Do not harvest the treated mature crop within 60 days after application. Consult the Attain* Herbicide Tank Mix labels for additional application instructions and use precautions.

NIMBLE herbicide + BANVEL* II Herbicide	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (excluding durum) and spring barley only, not underseeded to forages	
TIMING: Post-emergent application from the 2-5 leaf stage of spring wheat or spring barley and seedling stage to 8 leaf stage of kochia.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
Banvel* II Herbicide	0.110 litres/hectare
Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS	2 litres/1000 litres of spray solution
MIXING: Mixing order is NIMBLE herbicide then Banvel* II Herbicide then adjuvant. Refer to MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as kochia (including Group 2 resistant kochia).	
NOTES: Do not permit lactating dairy animals to graze fields within 7 days after application. Consult the Banvel* II Herbicide label for additional application instructions and use precautions	

NIMBLE herbicide + CURTAIL* M	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (excluding durum) and spring barley only	
TIMING: Post-emergent application from the full 3-leaf growth stage of spring wheat or spring barley.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
CURTAIL* M	1.5 litres/hectare
Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS	2 litres/1000 litres of spray solution
MIXING: Mixing order is NIMBLE herbicide , then Curtail* M, while maintaining agitation, then adjuvant, to the tank. Antifoaming agent may be required. Refer to MIXING DIRECTIONS section for general instructions	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as Canada thistle ¹ and wild buckwheat ² .	
Notes	
¹ Apply when the majority of thistles have emerged and are actively growing. For best control, apply before bud stage when thistles are no larger than 15 cm in height and before crop canopy prevents thorough coverage of thistles. Later emerging thistles will not be controlled.	
² Apply to actively growing wild buckwheat in the cotyledon to 5-leaf stage. Under environmental stress, such as dry weather, control may be reduced. Large plants may re-grow after treatment.	
NOTES: Consult the Curtail* M label for additional application instructions and use precautions	

NIMBLE herbicide + LONTREL* 360	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): Spring wheat (excluding durum) and spring barley only	
TIMING: Post-emergent application from the full 3-leaf to the flag leaf growth stage of spring wheat or spring barley.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
LONTREL* 360	0.21 litres/hectare
Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS	2 litres/1000 litres of spray solution
MIXING: Mixing order is NIMBLE herbicide , then Lontrel* 360, to the tank while maintaining agitation, then adjuvant. Antifoaming agent may be required. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as Canada thistle ¹ (seasonal control) and wild buckwheat ² .	
Notes	
¹ Apply when the majority of thistles have emerged and are actively growing. For best control, apply before bud stage when thistles are no larger than 15 cm in height and before crop canopy prevents thorough coverage of thistles. Later emerging thistles will not be controlled.	
² Apply to actively growing wild buckwheat in the cotyledon to 5-leaf stage. Under environmental stress, such as dry weather, control may be reduced. Large plants may re-grow after treatment	
NOTES: Consult the Lontrel* 360 label for additional application instructions and use precautions.	

NIMBLE herbicide + LONTREL* 360 + 2,4-D ESTER or MCPA ESTER	
LOCATION: Western Canada only.	
CROP(S): spring wheat (excluding durum) and spring barley only	
TIMING: Post-emergent application from full 3-leaf to just before flag leaf (shot blade) growth stage of crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
Lontrel 360	0.21 litres/hectare
Choose one of the following: 1) MCPA Ester (500 g/L) 2) 2,4-D Ester (600 g/L)	0.84 litres/hectare 0.7 litres/hectare
Adjuvants: AGRAL* 90, AG-SURF* or CITOWETT* PLUS	2 litres/1000 litres of spray solution
MIXING: Mixing order is NIMBLE herbicide first in fresh water, then Lontrel* 360 then 2,4-D Ester or MCPA Ester then adjuvant. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
All other weeds controlled by NIMBLE herbicide alone, as well as Canada thistle ¹ , lady's thumb, perennial sow thistle, stinkweed, volunteer canola, wild buckwheat and wild mustard.	
Notes	
¹ Provides seasonal control of Canada thistle.	
WEEDS SUPPRESSED	
All other weeds suppressed by NIMBLE herbicide alone, as well as cleavers.	
NOTES: Consult the Lontrel* 360 and MCPA Ester or 2,4-D Ester labels for additional application instructions, use Precautions and other label restrictions.	

SPRING WHEAT (INCLUDING DURUM)

NIMBLE herbicide + HORIZON* 240 EC	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only	
CROP(S): spring wheat (including durum) only	
TIMING: Post-emergent application from 2-leaf to flag leaf growth stage of the crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
HORIZON* 240 EC	230 mL/hectare for wild oats 290 mL/hectare for wild oats and green foxtail
Adjuvant: SCORE* only	800 mL/hectare for low rate of Horizon* 240 EC 1.0 litre/hectare for high rate of Horizon* 240 EC
MIXING: Mixing order is NIMBLE herbicide then Horizon* 240 EC then Score* adjuvant. DO NOT USE ADDITIONAL NON-IONIC SURFACTANT. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as wild oats and green foxtail.	
NOTES: Consult the Horizon* EC label for application instructions and use precautions.	

NIMBLE herbicide + MCPA ESTER 500 + HORIZON* 240 EC	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) only	
TIMING: Post-emergent application from 3-leaf to flag leaf growth stage of the crop. Apply to actively growing wild oats from the 1 to 6-leaf stage, prior to emergence of the 4th tiller and to green foxtail from the 1 to 5-leaf stage, prior to emergence of the 3rd tiller.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
MCPA ESTER 500	0.84-1.1 litres/hectare or 0.56 litres/hectare
HORIZON* 240 EC	230 mL/hectare
Adjuvant: SCORE* only	8 litres/1000 litres of spray solution (0.8% v/v)
MIXING: Mixing order is NIMBLE herbicide then Horizon* 240 EC then MCPA Ester 500 then Score* adjuvant. DO NOT USE ADDITIONAL NON-IONIC SURFACTANT. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Green foxtail ¹ Wild oats	
PLUS broadleaf weeds controlled or suppressed by NIMBLE herbicide and broadleaf weeds controlled by MCPA Ester 500 at 0.84-1.1 L per hectare or volunteer imazamox and imazethapyr tolerant canola (e.g. Clearfield* canola) (2-4 leaf stage) when MCPA is applied at 0.56 L per hectare.	
Notes	
¹ A reduction in control of green foxtail may be observed with the tankmix of NIMBLE herbicide + MCPA Ester 500 + Horizon* 240 EC.	
NOTES: Consult the MCPA Ester 500 and Horizon* 240 EC label for application instructions and use precautions.	

NIMBLE herbicide + ACCORD* + HORIZON* 240 EC	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (including durum) only	

TIMING: Make application from 2 to 5-leaf leaf growth stage of the crop. Apply to actively growing wild oats from the 1 to 6-leaf stage, prior to emergence of the 4 th tiller and to green foxtail from the 1 to 5-leaf stage, prior to emergence of the 3 rd tiller. Thorough coverage of foliage is important for consistent weed control.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
ACCORD* Herbicide	67 grams/hectare
HORIZON* 240 EC	230 mL/hectare
Adjuvant: SCORE* only	8 litres/1000 litres of spray solution (0.8% v/v)
MIXING: Mixing order is NIMBLE herbicide, then Accord then Horizon* 240 EC then Score* adjuvant. DO NOT USE ADDITIONAL NON-IONIC SURFACTANT. Refer to MIXING DIRECTIONS section for general instructions	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide, wild oats, green foxtail, and cleavers (1-4 whorl).	
NOTES: Prior to use, consult the Accord* label for additional rotational cropping restrictions, application instructions, use precautions, and other restrictions. Do not graze treated crops or cut for hay within 77 days of application.	

SPRING WHEAT (EXCLUDING DURUM)

NIMBLE herbicide + ACCORD* + AVENGE* 200-C	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (excluding durum) only NOT UNDERSEEDED TO FORAGES	
TIMING: Post-emergent application from 2 to 5-leaf growth stage of the crop.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare
ACCORD*	135 grams/hectare for most weeds listed below 165 grams/hectare for green foxtail
AVENGE* 200-C	3.5 liters/hectare
Adjuvant: MERGE* only	10 liters/1000 liters of spray solution (1.0% v/v)
MIXING: Mixing order is NIMBLE herbicide, then Accord*, then Avenge* 200-C while maintaining agitation, then MERGE* adjuvant. DO NOT USE ADDITIONAL NONIONIC SURFACTANT. Refer to MIXING DIRECTIONS section for general instructions.	
WEEDS CONTROLLED	
Weeds controlled or suppressed by NIMBLE herbicide alone and the NIMBLE herbicide + Accord* tank mix, as well as wild oats at populations between 1 and 200 plants per square meter	
NOTES: Do not apply this tank mixture more often than once every 2 years. Do not graze treated crops or cut for hay within 77 days of application. Consult the Accord* label for rotational cropping restrictions, application instructions, use precautions and other restrictions. Refer to the Accord* and Avenge* 200-C labels for additional information on appropriate weed staging and populations at application. Thorough coverage of foliage is important for consistent weed control. Avenge* 200-C can cause some crop injury. Always refer to the Accord* and Avenge* 200-C labels for additional application instructions, use precautions and restrictions.	

NIMBLE herbicide + HORIZON* 240 EC Herbicide + BANVEL* II Herbicide	
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.	
CROP(S): spring wheat (excluding durum) only, not underseeded to legumes	
TIMING: Post-emergent application from the 2-5 leaf stage of spring wheat and seedling stage to 8-leaf stage of kochia.	
TANK MIX COMPONENTS AND APPLICATION RATES	
NIMBLE herbicide	20 grams/hectare

HORIZON* 240 EC	0.230 L/Ha
Banvel* II Herbicide	0.110 L/Ha
Adjuvant: SCORE*	0.8 L/Ha
MIXING: Mixing order is NIMBLE herbicide then Banvel II Herbicide then Horizon 240EC then Score. Refer to MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha.	
WEEDS CONTROLLED	
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as kochia (including Group 2 resistant kochia), green foxtail and wild oats.	
NOTES: Do not permit lactating dairy animals to graze fields within 7 days after application. Consult the Banvel* II Herbicide and Horizon* 240 EC labels for additional application instructions and use precautions.	

NIMBLE herbicide + 2,4-D AMINE or ESTER + EVEREST* SOLUPAK* 70 DF or EVEREST* 70 DF							
LOCATION: For use in the prairie provinces and Peace River Region of British Columbia only.							
CROP(S): spring wheat (excluding durum) only NOT UNDERSEEDED TO LEGUMES							
TIMING: Post-emergent application from 1 to 6-leaf (1 to 4 leaves on main stem, plus 2 tillers) growth stage of the crop. For wild oats and green foxtail, apply from 1 to 6-leaf (1 to 4 leaves on main stem, plus 2 tillers)							
TANK MIX COMPONENTS AND APPLICATION RATES							
Component 1		Component 2		Component 3		Component 4	
NIMBLE herbicide	20 g/ha	Everest* 70 DF	43 g/ha			Agral*90 or Agsurf*	0.25% v/v
NIMBLE Herbicide	20 g/ha	Everest* 70 DF or Everest Solupak 70 DF	43 g/ha	2,4-D Amine or Ester	Up to 560 g ae/ha	Agral*90 or Agsurf*	0.25% v/v
MIXING: Mixing order is Everest* Solupak 70 DF or Everest* 70 DF then NIMBLE herbicide then 2,4-D Amine or Ester then adjuvant. Refer to MIXING DIRECTIONS section for general instructions.							
WEEDS CONTROLLED							
Broadleaf weeds controlled or suppressed by NIMBLE herbicide alone, as well as green foxtail and wild oats.							
NOTES: Observe minimum interval to harvest of 80 days after treatment. For black, dark brown and gray-wooded soil zones only, field peas may be grown the year following Everest* Solupak* application in fields where precipitation has been normal or above normal (10 year average) during the growing season, and where organic matter content is above 4% and pH below 7.5. Re-crop spring wheat only in brown soil zone. Consult the 2,4-D Amine or Ester and Everest* Solupak* 70 DF or Everest* 70 DF labels for additional information on rotational cropping restrictions, application instructions, use precautions and other restrictions.							

MINOR USES:

SEEDLING AND ESTABLISHED GRASS SPECIES FOR FORAGE AND SEED PRODUCTION

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The **DIRECTIONS FOR USE** for this product for the use described below were developed by persons other than Cheminova Canada and accepted for registration by Health Canada under

the User Requested Minor Use Label Expansion program. Cheminova Canada Company itself makes no representation or warranty with respect to performance (efficacy) or crop tolerance (phytotoxicity) claims for this product when used on the crops listed below. Accordingly, the User assumes all risks related to performance and crop tolerance arising, and agrees to hold Cheminova Canada Company harmless from any claims based on efficacy or phytotoxicity in connection with the use described below.

SEEDLING AND ESTABLISHED GRASS SPECIES FOR FORAGE AND SEED PRODUCTION

For broadleaf weed control, apply 20 grams of NIMBLE herbicide per hectare with AGRAL* 90, AG-SURF* or CITOWETT* PLUS at 2.0 litres per 1,000 litres of spray solution. (See "Mixing Instructions" section of the label).

Apply postemergence to the following Seedling or Established Grasses for Forage or Seed Production:

Meadow Bromegrass	Pubescent Wheatgrass
Smooth Bromegrass	
Tall Wheatgrass	Creeping Red Fescue
Slender Wheatgrass	
Tall Fescue	Streambank Wheatgrass
Orchard Grass	Northern Wheatgrass
Crested Wheatgrass	Western Wheatgrass
	Intermediate Wheatgrass

Apply postemergence to the following Established Grasses for Forage or Seed Production:

Kentucky Bluegrass

NIMBLE herbicide will control the broadleaf weeds listed on the label for **NIMBLE herbicide** alone. For best results apply to emerged, young, actively-growing weeds that are less than 10 cm tall or across. Thorough coverage of target weeds is essential. Apply with ground equipment only. Make only one application per growing season. The above grasses may not be grazed by or fed to livestock within 7 days of application. Please refer to the appropriate sections of this label for additional use precautions, application and/or mixing instructions.

MIXING INFORMATION

Mixing Precautions:

Do not allow the spray mixture to remain in the tank for more than 24 hours before spraying or the effectiveness may be reduced. If the spray mixture has been allowed to stand in the tank, use vigorous agitation to thoroughly disperse before resuming spraying.

Use the spray preparation of **NIMBLE herbicide** within 24 hours or product degradation may occur. Use spray preparation containing ASSERT* within 12 hours or product degradation may occur. If spray solution is left standing without agitation, thoroughly agitate and re-suspend any settled **NIMBLE herbicide** before resuming the spraying operation.

NOTE: Do not mix NIMBLE herbicide with liquid fertilizers, substances that contain boron or substances that release free chlorine. Mixing NIMBLE herbicide with any of these compounds will result in an insoluble substance in the tank.

Mixing Directions:

Select a spray volume that will ensure thorough coverage and a uniform spray pattern. Use a minimum of 55 L of spray volume per hectare. If tank mixing with other herbicides, always

consult the label of the tank mix partner(s) for minimum spray volume requirements and apply the tank mixture using a water volume recommended for all products.

1. Fill the spray tank $\frac{1}{4}$ to $\frac{1}{3}$ full with fresh water.
2. Add the recommended numberS/B amount of **NIMBLE herbicide** to the spray tank while the agitator is running. Continuous agitation is recommended through the entire mixing procedure to ensure products are completely dissolved and/or in suspension.
3. After **NIMBLE herbicide** has been fully dissolved and is in suspension, other tank mix components may be added while continuing to fill the tank with water. The proper sequence for tank mix components is:
 - a. dry formulations
 - b. emulsifiable concentrates
 - c. adjuvants (recommended adjuvant must be added to improve wetting and contact activity)
 - d. If a drift control agent is to be tank mixed, it is to be added first.
4. Refer to specific tank mixes in the DIRECTIONS FOR USE section of this label for mixing order and other mixing instructions.

For repeat loads, reduce the tank heel to 10% or less of the previous load. Fill tank with fresh water as described in step 1, and continue as directed. Emulsifiable concentrates may make dispersion of **NIMBLE herbicide** more difficult. For that reason, tank heels of the previous tank mix should be kept to 10% or less of the spray tank volume.

EQUIPMENT CLEANING INFORMATION

Cleaning Precautions:

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odor, which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR DESIRABLE VEGETATION.

Cleaning Instructions:

To avoid subsequent injury to crops other than wheat, spring barley or oats, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of **NIMBLE herbicide** from mixing and spray equipment as follows:

1. Drain tank and flush tank, boom and hoses with clean water for a minimum of ten minutes.
2. Visually inspect tank to ensure removal of all visible residues of **NIMBLE herbicide**. If necessary, repeat Step 1.
3. Fill the tank with clean water, then add 1 litre household AMMONIA (containing minimum of 3% ammonia) per 100 litres of water.
4. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation.
5. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.
6. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
7. Repeat Steps 3-5.
8. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **NIMBLE herbicide** is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to **NIMBLE herbicide** and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used

repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist.

Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the used of **NIMBLE herbicide** or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact Cheminova Canada at 1-888-316-6260 for further information.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

1. Make the empty container unsuitable for further use.
2. Dispose of the container in accordance with provincial requirements.
3. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

NIMBLE is a trademark of Cheminova A/S

*All other products mentioned are trademarks of their respective companies.

02NOV2009

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.