

PATROL® N/T

Liquid Inoculant Co-Pack for Soybean

Sufficient to Treat 1136 kg (2500 lbs. / 50 x 50 lb. units) of Seed

Manufactured for: United Agri Products Canada Inc.,
789 Donnybrook Drive, Dorchester, Ontario N0L 1G5
For General Inquiries : (800)-265-4624

Registration Number 2011011A Fertilizers Act

Net Weight: 3.2 kg

DIRECTIONS FOR IN-FURROW USE

Before using this product, read all the following instructions carefully.

PRIOR TO APPLICATION: Add the contents of the enclosed bottle of liquid *Bacillus subtilis* (100 mL) to the larger Rhizobium liquid inoculant bladder (3.1 L). Shake well to thoroughly mix.

APPLICATION RATE: 30 mL of Rhizobium/Bacillus mixture (30 g) per 304 meters of row.

The Rhizobium/Bacillus mixture must be diluted with cool, clean water and applied as indicated in the table below. Use the table to estimate application volumes for individual row widths.

ROW WIDTH	RECOMMENDED APPLICATION RATE*	CO-PACK UNIT TREATS
38 cm (15")	2640 mL/ha (1070 mL/acre)	1.2 ha (3 acres)
51 cm (20")	1960 mL/ha (790 mL/acre)	1.6 ha (4 acres)
76 cm (30")	1280 mL/ha (520 mL/acre)	2.5 ha (6 acres)
91 cm (36")	1130 mL/ha (460 mL/acre)	2.8 ha (7 acres)

The indicated volume of Rhizobium/Bacillus mixture should be diluted with water and applied at 45-90 litres per hectare for low volume sprayers, or 90-180 litres per hectare for larger sprayers. For a known row width, ensure the equipment is calibrated to deliver the correct application volume.

IN-FURROW APPLICATION

A. CLEAN. Ensure that the liquid application machinery is fully flushed of chemical residues before adding inoculant to spray tanks.

B. CONFIGURE. Configure the sprayer for in-furrow application. Select and install an appropriate flow regulator (orifice) disk in order to achieve a constant, uninterrupted stream of liquid into the furrow. Carefully adjust the position and flow of the liquid inoculant so that it falls directly onto the seed surface prior to covering in the furrow. Product efficacy may be limited if the inoculant is either applied above, or to the side of the seed in the furrow.

C. MIX. Prior to use, shake the product well in order to re-suspend any settled out components. Do not open the unit until required. Pour the complete contents into the spray tank. Adjust the volume of the spray tank to the required level using cool, dechlorinated water (**Refer to In-Furrow Application Table above**). Ensure that the inoculant is thoroughly mixed prior to use. To avoid any settling out of the nitrogen fixing bacteria, ensure that the spray tank volume is periodically agitated. The diluted inoculant must be used within 24 hours. Do not allow diluted tank mix to exceed 25°C.

D. SELECT RATE. The application rate per acre will vary according to row width. For guidance regarding specific application details and treatable areas **refer to In-Furrow Application Table above**.

DIRECTIONS FOR ON-SEED USE

Before using this product, read all the following instructions carefully.

PRIOR TO APPLICATION: Add the contents of the enclosed bottle of liquid *Bacillus subtilis* (100 mL) to the larger Rhizobium liquid inoculant bladder (3.1 L). Shake well to thoroughly mix.

- **PLANTER BOX APPLICATION:** Shake the Rhizobium / Bacillus mixture well, and apply to the seed at the rate of 77 mL of inoculant (77 g) per 27.2 kg (1 bushel) of seed.
- **BULK SEED APPLICATION:**
 1. Remove cap. Insert gasket into valve body. Attach flow valve and hose to bladder.
 2. Invert bladder above treatment area so the end of the hose is just above the seed (for accurate application rates, ensure hose is straight when dispensing inoculant).
 3. Adjust flow valve to match output of grain auger. (See table below).
 4. To ensure adequate mixing of seed and inoculant it is recommended to run auger at 50% capacity.
 5. Assess the application rate several times during inoculation to ensure correct target flow rate.

Plant all treated seed as soon as possible, and within a maximum of 10 days following inoculation. Keep treated seed as cool as possible (1 - 20°C).

SEED TREATMENT COMPATIBILITY

Some seed treatments are harmful to liquid inoculants. Contact your dealer /manufacturer for current details. Please observe planting window recommendations for optimal responses.

Flow Valve Setting	Inoculant Flow Rate	Seed/Auger Flow Rate		
	mL/min	kg/min	Lbs/min	Approx. bu (50lbs units)/min
1	360	127	279	5 (6)
2	860	303	667	11 (13)
3	1340	473	1041	17 (21)
4	1660	586	1289	22 (26)
5	1780	628	1382	23 (28)
6	2030	717	1577	26 (32)

GUARANTEE

This product contains a minimum of 3.0×10^9 viable cells of *Bradyrhizobium japonicum* per gram and 2.2×10^{10} viable cells of *Bacillus subtilis* per gram.

STORAGE/PRECAUTIONS

Store below 20°C. Do not allow product to freeze. Ensure that inoculant is stored correctly in the field prior to use. Once opened, use within 24 hours. Plant seed as soon as possible after inoculation and within the maximum stated period above. Do not use inoculant that is past its expiry date or that has not been correctly stored. Unless otherwise directed, do not mix inoculant with caustic fertilizers or pesticides. If soils are especially dry at planting, any risk of damage to the inoculant can be minimized through light irrigation. Do not feed seed treated with this product to livestock. Store away from food or feed. Avoid prolonged or repeated exposure to skin. See material safety data sheet for additional information.

IN CASE OF EMERGENCY DUE TO A MAJOR SPILL, FIRE OR POISONING INVOLVING THIS PRODUCT CALL DAY OR NIGHT, 1-800-561-8273.

United Agri Products Canada Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purposes stated on such label only when used in accordance with the directions and under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of the product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of United Agri Products Canada Inc. In no case shall United Agri Products Canada Inc. be liable for consequential, special or indirect damages (including without limitation loss of profits or revenues) resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Except or expressly provided herein, United Agri Products Canada Inc. makes no warranties, guarantees, conditions or representations of any kind, either

express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. Buyer's or user's exclusive remedy, and United Agri Products Canada Inc. total liability, shall be for damages not exceeding the cost of the product.

Patrol is a registered trademark of United Agri Products Canada Inc.