

Penncozeb® 75DF

Fungicide

For excellent control of early and late blight in potatoes.

Penncozeb® fungicide delivers...

- Effective broad-spectrum multi-site control with a proven history of success
- Convenience for you - provides a wide window of application, protection from planting through harvest
- Rotation and tank mix options with systemic and other protectant fungicides
- A great tool and ideal partner for your disease resistance management program
- Application flexibility allows you to choose how to apply - ground or air



Application Recommendations for Potatoes

Begin applications early

- Apply 1.1 kg per hectare when plants are 10 – 15 cm high
- Increase to 1.75 kg per hectare as plants increase in size
- Increase to 2.25 kg per hectare at row closure
- Apply every 7 – 10 days throughout the season
- During periods of wet weather favoring late blight and/or vigorous crop growth, spray intervals may be reduced to 5 - 6 days
- Apply 1.1 kg per hectare until row closure, then increase the rate to 1.75 kg per hectare until conditions allow the return to the regular spray schedule
- Thorough, uniform coverage is essential for good disease control
- Use 40 liters of water per hectare for aerial applications
- Do not apply within 1 day of harvest



West: 1-800-561-5444
Ontario and Maritimes:
1-800-265-5444
Quebec: 1-800-361-9369
www.uap.ca



See product label for specific use instructions. Always read and follow label directions and precautions. Penncozeb is a registered trademark of United Phosphorus, Inc. © June 2008 United Phosphorus, Inc. www.upi-usa.com