



SERENADE[®] MAX – For Proven Control of Bacterial Disease on Fruit & Vegetables




Control bacterial diseases in fruits and vegetables with the proven power of SERENADE[®] MAX. Powered by the complex biochemistry of *Bacillus subtilis* (QST 713), SERENADE MAX effectively attacks spores and prevents infection. SERENADE MAX is safe for use on fresh &

processed fruit and vegetables and features a unique, multi-site mode of action that makes it the perfect weapon to combat quickly mutating disease pathogens, delaying resistance. Improve your disease control program in fruits and vegetables with SERENADE MAX.



CROP	DISEASE	RATE (kg/ha)	Application Instructions
Asparagus	Botrytis blight	3,0-6,0	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.



Celery	Pink Rot	3,0-6,0	Make the first application approximately 8 weeks before harvest and repeat on a 14-day interval. Apply Serenade MAX as a directed spray in sufficient water to ensure thorough coverage of the base of the plants and the surrounding soil surface.
--------	----------	---------	---



Bulb Vegetables Onions (including green bunching, dry bulb and Welch) Garlic (including Elephant garlic), leeks, shallots.	Botrytis neck rot Botrytis leaf blight	3,0-4,5	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.
	Downy mildew	3,0-6,0	



Cole Crops Broccoli, Brussels sprouts, Cabbage, Cauliflower, Cavalo Broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard Spinach, Rape Greens.	Watery Soft Spot	3,0-6,0	Make the first application at emergence, or immediately following transplanting. Repeat applications on 10-14 day intervals if conditions for disease development continues.
--	------------------	---------	--



Cucurbits Cucumber, Cantaloupe, Melon, Muskmelon, Squash, Watermelon, Pumpkin.	Powdery Mildew	3,0-6,0	Begin application soon after emergence or transplant, when conditions are conducive for disease development. Repeat applications on an interval of 7-10 days.
---	----------------	---------	---



 CANADA



CROP	DISEASE	RATE (kg/ha)	Application Instructions
Fruiting Vegetables Eggplant, Ground Cherry Pepino, Peppers (all varieties), Tomatillo and Tomatoes	Early Blight	4,5	Make the first application when plants are 6-10 cm high, or when conditions are conducive for disease development. Repeat applications on an interval of 5-7 days.
	Botrytis Gray Mould	3,0-6,0	
	Powdery Mildew ***	3,0-6,0	Begin application soon after emergence or transplant, and repeat on an interval of 7-10 days.
			* For use in Tomatoes and Peppers only



Legume Vegetables Beans (adzuki, broad, dry, field, kidney, lablab, lima, moth, mung, navy, pink, pinto, rice, runner, snap, tepary, urd, wax, yardlong, catjang, chick pea, cowpea, guar, jackbean) Peas (black-eyed, crowder, dwarf, edible podded, English, field, garden, green, pigeon, snow, southern, succulent, sugar snap) Lentils	White Mould	3,0-6,0	Make the first application at planting (or immediately following planting but prior to crop emergence). Make a second application as a directed spray with multiple nozzles per seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning. Repeat applications on 7-14 day intervals if conditions for disease development persists.
	Botrytis pod rot	3,0-6,0	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.



Lettuce	Downy mildew Powdery mildew	3,0-6,0	Begin applications at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.
	Sclerotinia head and leaf drop	3,0-6,0	Make the first application at planting (or immediately following planting but prior to crop emergence). Make a second application as a directed spray with multiple nozzles per seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning. Repeat applications on 10-14 day intervals if conditions for disease development persist.



Strawberry	Botrytis gray mould	3,0-6,0	Begin application at the first sign of disease, or when conditions become conducive for disease development. Repeat as necessary on a 7-10 day interval.
------------	---------------------	---------	--

CONTACT: