

Supra Captan 80 WDG

FUNGICIDE

Supra Captan 80 WDG is a water dispersible granular fungicide for control of a wide range of diseases in fruits, vegetables, ornamentals, golf courses and sod farms. Supra Captan 80 WDG is a contact and protective fungicide formulated for dust-free application.

ACTIVE INGREDIENT: Captan 80%.

CROPS: Apples, apricot, blackberry, blueberry, cucumber, grape, loganberry, nectarine, peach, pear, plum, prune plum, raspberry, rhubarb, sour and sweet cherry, strawberry, tomato, golf courses and sod farms, soil and greenhouse bench treatments for seedling and transplants, tobacco seedlings, and many ornamental flowers.

RATE/TIMING: 0.6-6.0 kg/ha (0.24-2.4 kg/acre).
75-156 g/1000 square metres for greenhouse applications. Rates and timing depends on crop and targeted plant disease. Except as specified, begin application before or at the first sign of disease and repeat as necessary to maintain control. Consult label for complete directions. Do not apply immediately before or closely after oil sprays.

WATER VOLUME: For dilute ground applications, use recommended rate of Supra Captan 80 WDG in 1000 L of water unless otherwise specified on the label. For low volume sprayer applications, maintain the Supra Captan rate/ha as if spraying with a higher water volume.

DISEASES CONTROLLED

Anthracnose	Fruit rot
Bitter rot	Grey mold rot
Black knot	Late blight
Black rot	Leaf spot
Black spot	Melting-out
Botrytis flower blight	Mummy berry
Brooks fruit spot	Petal blight
Brown patch	Pythium
Brown rot	Root rot
Bulb rots	Scab
Cane spot	Septoria leaf spot
Damping-off and fungus root diseases	Shot hole (leaf spot)
Dead arm	Sooty blotch
Early blight	Spur blight
Fly speck	



QUICK SHEET

ALWAYS READ THE LABEL BEFORE USE

TANK MIXES: Supra Captan 80 WDG can be used safely and effectively at recommended rates with most of the commonly used insecticides and fungicides with the exception of oil and strongly alkaline materials, including calcium arsenate.

CAUTION: Do not use on D'Anjou pears. Do not apply in combination with or immediately before or closely following oil sprays. Do not combine with emulsifiable liquids in the same spray tank unless compatibility has been proven. Combination of Supra Captan 80 WDG and sulphur should not be used on sulphur sensitive crops. Do not use on home lawns, parks, schoolyards, or other recreational areas.

RAINFAST: Once dried on crop.

PREHARVEST INTERVAL: 2 days for most crops except 5 days for sour cherries and 7 days for apples, grapes and pears.

RE-ENTRY: Once dried on crop.

PACKAGE SIZE: 2 x 5 Kg.

ADDITIONAL: Used at high rates or in drenching sprays, Supra Captan 80 WDG may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plum and cherries. To avoid leaf spotting under warm, cloudy, humid conditions, use at lowest recommended rates and avoid drenching trees.

SURFACTANT: The use of LI700 to lower the pH of the spray water prior to adding Supra Captan 80 WDG can assist in product penetration into the leaf surface. The use of spreaders that cause excessive wetting is not advised.

RE-CROPPING: No restrictions.

STORAGE: Store in a cool, dry, ventilated place, away from feeds and foods. Keep away from heat, fire and sparks.

GROUP: (m) multi-site activity.

NOTES:



West: 1-800-561-5444 Ontario: 1-800-265-4624 Quebec & Maritimes: 1-800-361-9369