

# Folpan® 80WDG

## FUNGICIDE

Folpan 80WDG is a water dispersible granular fungicide for the control of a number of fungal diseases on certain fruit, vegetable, ornamental and other crops.

**CAUTIONS:** Fungicidal activity may be reduced with strongly alkaline materials such as hydrated lime. Do not use in combination with or closely following oil sprays. Keep agitation running, do not allow the mixture to stand without agitation. Do not use in low volume, gear-type application equipment. Do not apply by aerial application.

Do not apply by chemigation or through any type of irrigation system. For commercial use only. Do not use in and around home or residential areas such as parks, schools, public buildings, playing fields or any other areas where the general public including children could be exposed.

**PACKAGE SIZE:** 2 x 5 kg.

**RE-ENTRY:** 24 hours.

### FRUIT CROPS

Crop	Disease/Rate	Timing	PHI*
Apples & crabapples	Alternaria leaf spot, black rot, Brooks spot, fly-speck, scab, sooty blotch: 3.0-3.75 kg/ha in sufficient water for good coverage (up to 3000 L/ha).	Green tip until harvest (in western Canada do not apply between bloom and July to avoid injury). Russetting may occur in Eastern Canada if applied between bloom and 30 days after petal fall on some varieties. Do not apply more than 6 times per season.	1 Day
Cranberries	Fruit rot: 3.5 kg/ha use up to 2000 L/ha.	5% blossoms open and then 10-14 days later when 50% -75% blossoms open. Maximum of 2 sprays per season.	30 Days
Grapes	Dead arm: 1.25 kg/ha in enough water for good coverage (up to 1000 L/ha).	New shoots 2.5-5.0 cm long. Repeat when growth 10-15 cm long. Do not apply more than 2 applications per season.	1 Day
	Black rot, downy mildew: 1.25 kg/ha in enough water for good coverage (up to 1000 L/ha).	Just before bloom, just after bloom and at 1 <sup>st</sup> cover spray. 2-3 weeks later for powdery mildew. Apply no more than 4 times per season.	1 Day
Strawberry	Gray mold, fruit rot, leafspot: 2.5 kg/ha. Use up to 2000 L/ha to obtain adequate coverage.	Before 1 <sup>st</sup> infection, at 7 day intervals until harvest. Do not apply more than 6 times per season.	1 Day

\*Pre Harvest Interval



## QUICK SHEET

## ALWAYS READ THE LABEL BEFORE USE

VEGETABLE CROPS			
Crop	Disease/Rate	Timing	PHI*
Field Cucumbers, melons, pumpkin, & squash	Anthracoise & downy mildew 2.5-5 kg/ha. Use up to 2000 L/ha (enough for thorough coverage).	Apply when 1 <sup>st</sup> true leaves appear. Repeat at 7 day intervals until harvest. Maximum of 6 applications per season.	1 Day
Field Tomatoes	Anthracoise 5 kg/ha. Use up to 2000 L spray solution/ha.	During 1 <sup>st</sup> bloom and repeat at 7 day intervals until harvest. Do not apply more than 6 times per season.	1 Day

\*Pre Harvest Interval

ORNAMENTAL CROPS		
Crop	Disease/Rate	Timing
Azaleas	Stem root rot of cuttings: 1.0-1.9 kg/1000 L.	Soak cutting for 15-30 minutes before planting.
Carnations	Blight (Alternaria leaf spot): 1.25 kg/1000 L.	2 weeks after planting and 3-4 times at bi-weekly intervals. Maximum of 6 sprays per season.
Chrysanthemums	Septoria leaf spot: 1.25 kg/1000 L.	Immediately after planting 7-10 day intervals. Do not apply more than 6 times per season.
Iris	Didymellina leaf spot: 1.25 kg/1000 L.	When plants emerge and repeat at 7-10 day intervals. No more than 4 applications per season.
Marigolds, Zinnias	Alternaria Leaf Spot: 1.25 kg/1000 L.	Apply when plants emerge and repeat at 3-7 day intervals depending on growth rate. Do not apply more than 6 times per season.
Poinsettias (Greenhouse)	Pythium root rot: 1.25-1.4 kg/1000 L 0.57 L/ 20 cm pan.	Apply at planting time and repeat 10 days later. Maximum of 2 applications per season.
Snapdragon	Anthracoise: 1.25 kg/1000 L.	When plants emerge and repeat at 3-7 day intervals. No more than 6 applications per season.

**RESIDUAL CONTROL:** 3-10 days depending on the rate of growth in the crop.

**ADDITIONAL:** May be combined with most commonly used insecticides, adjuvants, and fungicides, however check compatibilities of tank mixes before using. Do not allow solutions to stand without agitation.

**ACTIVE INGREDIENT:** Folpet 80%.

**STORAGE:** Store in cool, dry, ventilated place away from feed and foods.

**GROUP:** M Fungicide.



[www.uap.ca](http://www.uap.ca)

**West: 1-800-561-5444 Ontario & Maritimes: 1-800-265-5444 Quebec: 1-800-361-9369**