

# UP-Cyde<sup>®</sup> 2.5 EC

## Insecticide

***For broad-spectrum pest control on numerous vegetables, including asparagus, carrots, celery, cole crops, corn, lettuce, onions, potatoes, rutabagas, turnips and tomatoes.***

### UP-Cyde Use Rates for Vegetables

<i>Vegetable</i>	<i>Pests</i>	<i>Application</i>
Asparagus	Asparagus Beetle	Apply 140 mL/ha to spears & ferns when insects are observed. Repeat application as required.
Carrots	Carrot Rust Fly	Apply 280 mL/ha with 550 L of water when insects are first noticed. Do not apply more than 3 times per season or 35 days before harvest.
Celery	Potato Leafhopper	Apply 140 mL/ha with 500 L of water when insects are first noticed.
Cole Crops (Cabbage, Cauliflower, Broccoli, Brussel Sprouts)	Cabbage Looper, Imported Cabbage Worm, Diamond-back Moth Larvae	Apply 140 mL/ha & add AGRAL 90 surfactant at 0.03% v/v (300 mL / 1000 L of spray mix). Use as needed at 2 week intervals but not more than 3 times per season, or three days before harvest.
	Thrips	Apply 200 mL/ha & add AGRAL 90 surfactant at 0.03% v/v (300 mL / 1000 L of spray mix).

## UP-Cyde Use Rates for Vegetables

<i>Vegetable</i>	<i>Pests</i>	<i>Application</i>
Corn (Sweet & Field)	European Corn Borer	Apply 280 mL/ha in 325 to 450 L of water. Spray no later than when first damage appears & repeat at 4-7 day intervals.
Lettuce	Aster Leafhopper	Apply 200 mL/ha when insects or damage first appear. Apply up to 4 times per season at 7-14 day intervals & not 14 days before harvest.
Onions	Onion Maggot	Apply 280 mL/ha later in the growing season or on windrow onions prior to harvest.
	Thrips	Apply 280 mL/ha at 10 day intervals but not more than 3 times per season or 3 days before harvest.
Potatoes	Colorado Potato Bee- tle, Potato Flea Beetle, Potato Leafhopper, Tuber Flea Beetle	Apply 140 mL/ha when insects or damage first appear.
	Tarnished Plant Bugs	Apply 200 mL/ha when insects appear.
Rutaba- gas & Turnips	Crucifer Flea Beetle	Apply 200 mL/ha when insects or damage first appear. Do not apply 21 days before harvest.
Tomatoes	Colorado Po- tato Beetle, Potato Flea Beetle, Potato Leafhopper	Apply 140 mL/ha when insects or damage first appear. Repeat as required, but not 3 days before harvest.



**UPI**

*Built for where plant protection is going*

Always read and follow label directions and precautions. UP-Cyde is a registered trademark of United Phosphorus, Inc. © January 2008 United Phosphorus, Inc. UCYD-805