

UBG

**ZINEB 80 W**

COMMERCIAL

CAUTION

POISON

READ THE LABEL BEFORE USING

REGISTRATION NO. 9318

PEST CONTROL PRODUCTS ACT

GUARANTEE: .....zineb 80%

UNITED AGRI PRODUCTS CANADA INC.

789 Donnybrook Drive

Dorchester, Ontario

N0L 1G5

6842 01/00

NET: 10 kg

**DIRECTIONS FOR USE:**

**MIXING METHOD:** Add half of the required amount of water to the spray tank; agitate continuously. Add the prescribed amount of ZINEB 80 W by flushing it through a screen with a hose; or premix Zineb into a slurry in a separate container and add this mixture to a nearly filled spray tank. Add the balance of water. Do not allow the spray mixture to stand.

This product is compatible with most insecticides and fungicides. Do not combine wettable powders with emulsifiable liquids in the same spray tank. Always use sufficient water (1000-1500 L per ha) to ensure proper coverage of foliage. Apply when disease first appears. Use the lower rate early in the season and increase the rate as the plants grow. Do not apply before harvest within the number of days shown after each crop.

CROP	DISEASE	APPLICATION	Days to Harvest
ASPARAGUS	Rust	1.1-3.3 kg/ha; 5-10 day intervals. Do not apply to spears which are to be cut.	
BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER	Alternaria leaf spot, Downy mildew	1.1-3.3 kg/ha; 5-10 day intervals.	7 days
CARROTS	Alternaria blight, Cerospora blight	1.1-3.3 kg/ha; 7-10 day intervals	7 days
CELERY	Early blight, Late blight	1.1-3.3 kg/ha; 3-5 day intervals when seedlings first emerge. Wash and trim.	14 days
CUCUMBERS, MELONS, PUMPKINS, SQUASH	Alternaria leaf spot, Anthracnose, Downy mildew, Gummy stem blight, Septoria leaf spot	2.3-2.8 kg/ha; 7-10 day intervals when plants begin to vine or when disease threatens.	5 days
EGGPLANT, PEPPERS	Anthracnose, Early blight, Late blight	1.1-3.3 kg/ha; 7-10 day intervals.	5 days
LETTUCE	Downy mildew	1.7-2.3 kg/ha; 7-10 day intervals.	10 days
ONIONS	Botrytis leaf blight, Downy mildew, Purple blotch	0.85-3.3 kg/ha; 7-10 day intervals.	7 days
POTATOES	Early blight, Late blight	1.7-3.3 kg/ha; 7-10 day intervals.	1 day
TOMATOES	Anthracnose, Grey mold, Early blight, Late blight, Grey leaf spot, Septoria leaf spot	0.85-3.3 kg/ha; 7-10 day intervals.	5 days

**PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.** May cause irritation of nose and throat. Avoid inhalation of spray or dust, and contact with skin, eyes, and clothing. Do not contaminate food or feed. Do not contaminate ponds, lakes or streams. Keep container tightly closed when not in use. Keep away from fire and sparks. If allowed to become wet or overheated in storage, chemical changes may occur which will impair fungicidal effectiveness, and may also generate flammable vapours.

**FIRST AID:** If on skin, wash with plenty of water. If in eyes, flush with plenty of water for 15 minutes and get medical attention. Take container, label or product name and Pest Control Product Registration number with you when seeking medical attention

**DISPOSAL:**

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills .

**NOTICE TO USER:** This control product is to be used in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

**NOTICE TO BUYER:** Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.