

# Sword®

## HERBICIDE

For systemic control of many broadleaf weeds in: barley, canary seed, oats, summerfallow, wheat, and seedling and established grasses grown for forage.

### WEEDS CONTROLLED:

Buckwheat (wild, tartary, volunteer cultivated)	Mustard (wild, ball, tall, wormseed, yellow)
Cleavers*	Night flowering catchfly
Common ragweed	Pigweed (prostrate, redroot)
Corn spurry	Russian thistle, ***
Cow cockle	Shepherd's purse
Flixweed	Annual sow-thistle,
Green smartweed	Stinkweed
Hemp-nettle**	Volunteer rapeseed
Knotweed	Volunteer sunflower
Kochia	Also chickweed, wild oats, and millet by tank mixes.
Lady's-thumb	
Lamb's-quarters	

### TOP GROWTH CONTROL ONLY:

Canada thistle	Hedge bindweed
Field bindweed	Perennial sow-thistle

\* Cleavers: spray in 1-2 whorl stage.

\*\* Hemp-nettle: spray before second pair of true leaves.

\*\*\* Russian thistle: spray when less than 5 cm in height.

**CROPS:** Barley, canary seed, durum wheat, oats, summerfallow, wheat, winter wheat, and seedling and established grasses grown for forage. (Not underseeded to legumes.)

**RATE:** Apply 1.0-1.5 L/ha (.4-.6 L/acre) (one 10 L jug treats 25 to 16.7 acres).

**WATER VOLUME:** Use 50-100 L/ha (5-20 gal/acre) by ground and minimum of 30 L/ha (3 gal/acre) by air.

**TANK MIXES:** Horizon®, Everest, (wheat only). Nortrace plant nutrients. (Conduct a jar test with Nortrace plant nutrients before tank mixing.) For other tank mixes see label.

### TIMING:

**Barley:** 2 to 4 leaf.

**Wheat:** (spring, durum), oats, canary seed, 2 to 5 leaf.

**Winter Wheat:** apply in the spring before the crop is more than 30 cm high (top leaf extended).

**WINTER ANNUAL WEEDS:** Should be sprayed as soon as new growth starts.

**CAUTION:** Best results are obtained when weeds are in the 2-3 leaf stage and actively growing. Less than satisfactory control may result if weeds are not actively growing.



